

# Northumbria Research Link

Citation: Cellier-Rastit, Marie, James, Arthur L., Orenge, Sylvain, Perry, John D., Robinson, Shaun N., Turnbull, Graeme and Stanforth, Stephen P. (2019) Fluorogenic L-alanylaminopeptidase substrates derived from 6-amino-2-hetarylquinolines and 7-amino-3-hetarylcoumarins and their potential applications in diagnostic microbiology. *Bioorganic & Medicinal Chemistry Letters*, 29 (10). pp. 1227-1231. ISSN 0960-894X

Published by: Elsevier

URL: <https://doi.org/10.1016/j.bmcl.2019.03.011>  
<<https://doi.org/10.1016/j.bmcl.2019.03.011>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/38383/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

# Fluorogenic L-alanylaminopeptidase substrates derived from 6-amino-2-hetarylquinolines and 7-amino-3-hetarylcoumarins and their potential applications in diagnostic microbiology

Marie Cellier-Rastit,<sup>a</sup> Arthur L. James,<sup>b,d</sup> Sylvain Orenge,<sup>a</sup> John D. Perry,<sup>c</sup> Shaun N. Robinson,<sup>b</sup> Graeme Turnbull<sup>b</sup> and Stephen P. Stanforth<sup>b\*</sup>

<sup>a</sup> Research & Development Microbiology, bioMérieux SA, 3 route de Port Michaud, 38 390 La-Balme-les-Grottes, France

<sup>b</sup> Department of Applied Sciences, Northumbria University, Newcastle upon Tyne, NE1 8ST, UK

<sup>c</sup> Department of Microbiology, Freeman Hospital, Newcastle upon Tyne, NE7 7DN, UK


<sup>d</sup> deceased May 2014

\*Corresponding authors. Tel.: +44-191 227 4784; fax: + 44-191 227 3519; e-mail: steven.stanforth@northumbria.ac.uk

**Abstract.** Six novel fluorogenic enzyme substrates for detecting L-alanylaminopeptidase activity in microorganisms have been prepared and evaluated in Columbia agar media. The substrates are L-alanyl derivatives of 6-amino-2-hetarylquinolines and 7-amino-3-hetarylcoumarins. Both the quinoline and coumarin series of substrates produced fluorescence in the presence of Gram-negative microorganisms. In contrast, fluorescence generation in the presence of the Gram-positive microorganisms and yeasts was limited or absent.


**Keywords:** Fluorogenic substrates, L-alanylaminopeptidase, microorganism detection, quinolines, coumarins

The detection of L-alanylaminopeptidase activity in microorganisms often provides a useful insight as to whether a microorganism of interest can be categorised as either Gram-negative or Gram-positive. This is because Gram-negative microorganisms are usually associated with comparatively high levels of L-alanylaminopeptidase activity compared to their Gram-positive counterparts.<sup>1,2</sup> One strategy for detection of L-alanylaminopeptidase activity involves monitoring the increase in fluorescence when a weakly fluorescent substrate **1** is converted into a highly fluorescent heterocyclic amine derivative **2** as depicted by the enzymatic transformation shown in Scheme 1.




**Scheme 1.** Generation of fluorescence resulting from L-alanylaminopeptidase activity (Het = heteroaromatic).

We have previously reported the synthesis and evaluation of fluorogenic enzyme substrates **3** (X = O, S),<sup>3</sup> and derivatives of these compounds, for the detection of L-alanylaminopeptidase activity in a selection of clinically important microorganisms in Columbia agar media (Figure 1). Good growth of the Gram-negative microorganisms was observed on this medium and, as anticipated, fluorescent colonies were produced by hydrolysis of these substrates. In comparison, the growth of the Gram-positive microorganisms on the medium was either reduced or even inhibited by the substrate, and consequently only weakly fluorescent colonies were produced where microorganism growth had occurred. The fluorogenic substrate **4** was also prepared but, disappointingly, it only gave moderately fluorescent colonies with Gram-negative microorganisms.<sup>4</sup> Thus, we were interested in preparing and assessing whether heterocyclic systems related to structure **4** would produce more intensely fluorescent colonies with Gram-negative microorganisms. The quinoline derivatives **5a-c** and the coumarin derivatives **6-8**, in which the central 1,4-disubstituted phenyl ring in structure **4** is replaced by a 2,6-disubstituted quinoline ring and a 3,7-disubstituted coumarin ring respectively, were therefore chosen for study as potential substrates for the detection of L-alanylaminopeptidase activity in microorganisms. It was anticipated that hydrolysis of these substrates would produce the corresponding fluorescent amine derivatives **9a-c** (from quinolines **5a-c** respectively) and amines **10-12** (from the coumarins **6-8** respectively) (Figure 2). An *N,N*-disubstituted derivative of the amine **9c** has been previously incorporated into a two-photon fluorescent probe for nitric oxide detection in cells and tissues.<sup>5</sup> Related to amines **9**, are derivatives of 2-amino-6-hetarylnaphthalenes (general structure **9**, quinoline nitrogen = CH) which have been prepared and derivatised for a variety of fluorescence applications.<sup>6</sup> 7-Aminocoumarin derivatives are well known fluorescent heterocycles e.g. the L-alanyl derivatives of 7-amino-4-methylcoumarin<sup>7</sup> and 7-amino-4-trifluoromethylcoumarin<sup>8</sup> are substrates that have been utilised for the detection of L-alanylaminopeptidase activity. The aminocoumarin **10** has been described in the literature as an intermediate in the preparation of the corresponding isothiocyanate (**10**, NH<sub>2</sub> = NCS).<sup>9</sup>




**Figure 1.** Fluorogenic L-alanylaminopeptidase substrates.



**Figure 2.** Fluorescent amines **9a-c** and **10-12** resulting from the hydrolysis of substrates **5a-c** and **6-8** respectively.


The synthesis of the required series of substrates **5a-c** is depicted in Scheme 2. 6-Nitroquinoline-2-carboxaldehyde **13**<sup>10</sup> was treated with *N*-methyl-1,2-phenylenediamine in the presence of Oxone<sup>®</sup> giving the nitroquinoline derivative **16a** via oxidation of the initially formed dihydro-derivative **14a**. Heterocycle **13** and 2-aminothiophenol in hot dimethylsulfoxide solution produced the benzothiazole derivative **16c** via the dihydro intermediate **14c**. The reaction of aldehyde **13** with 2-aminophenol gave a Schiff base **15b** which was subjected to an oxidative cyclisation when treated with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) producing the benzoxazole derivative **16b**. All three nitroquinolines **16a-c** were reduced with tin(II) chloride affording the amines **9a-c** respectively. These amines were then reacted with Boc-L-alanine in a mixed anhydride condensation reaction and the resulting the Boc-protected amides **17a-c** were deprotected using a saturated solution of hydrogen chloride in ethyl acetate furnishing the required substrates **5a-c**.



**Scheme 2.** Synthesis of the substrates **5a-c**. Reagents and conditions: (i) *N*-methyl-1,2-phenylenediamine, Oxone<sup>®</sup>, DMF/H<sub>2</sub>O (25:1), rt, 16 h; (ii) 2-aminothiophenol, DMSO, 110 °C, 16 h; (iii) 2-aminophenol, EtOH, reflux, 16 h; (iv) DDQ, CH<sub>2</sub>Cl<sub>2</sub>, rt, 16 h; (v) SnCl<sub>2</sub>·2H<sub>2</sub>O, EtOH, reflux, 16 h; (vi) Boc-L-alanine, *N*-methylmorpholine, isobutyl chloroformate, THF, -5 °C then add **9a-c**; (vii) sat. HCl/EtOAc, rt, 2 h.

The synthetic route adopted for the preparation of substrates **6-8** is outlined in Scheme 3. 3-Aminophenol **18** was subjected to a mixed anhydride coupling reaction with Boc-L-alanine giving the phenol derivative **19**. Compound **19** was then reacted with paraformaldehyde in the presence of MgCl<sub>2</sub> under basic conditions<sup>11</sup> producing the salicylaldehyde derivative **20**. The reaction of compound **20** with the appropriate ethyl ester derivatives of hetarylacetic acids under basic conditions afforded the hetarylcoumarins **21** and **22**. Treatment of coumarin **22** with methyl iodide furnished the pyridinium salt **23**. Removal of the Boc protecting-group from compounds **21-23** was

achieved by treatment with a saturated solution of hydrogen chloride in ethyl acetate giving the required substrates **6-8**.



**Scheme 3.** Synthesis of the substrates **6-8**. Reagents and conditions: (i) Boc-L-alanine, *N*-methylmorpholine, isobutyl chloroformate, THF, -5 °C then add **18**; (ii) paraformaldehyde, Et<sub>3</sub>N, MgCl<sub>2</sub>, CH<sub>3</sub>CN, reflux, 48 h; (iii) piperidine (0.6 equiv.), EtOH, reflux, 18 h; (iv) sat. HCl/EtOAc, rt, 2 h; (v) MeI, CH<sub>3</sub>CN, rt, 24 h.

Each of the substrates **5a-c** and **6-8** were evaluated in Columbia agar medium (37 °C in air for 18 hours) against a panel of 20 clinically important microorganisms, including 10 Gram-negative bacteria, 8 Gram-positive bacteria and 2 yeasts. Columbia agar media was selected because each substrate could be evaluated simultaneously against the complete panel of 20 microorganisms on a single plate. The growth of the microorganisms was compared to control plates in which no substrate was present. The Gram-negative microorganisms all exhibited strong growth on the control plates whereas the Gram-positive microorganisms and the yeasts showed only moderate growth.

The fluorescence produced from the hydrolysis of the L-alanylaminopeptidase substrates **5a-c** by the panel of 20 microorganisms is shown in Table 1 and Figure 3 shows a photograph of a representative agar plate depicting the fluorescence produced from substrate **5c**. There was good growth of all the Gram-negative microorganisms in the presence of substrates **5a-c** and strongly fluorescent colonies were observed with nearly all these bacteria. This activity profile is similar to that observed with the commercially available substrate, L-alanyl-7-amino-4-methylcoumarin (data not shown). Growth of

the Gram-positive microorganisms and yeasts in the presence of substrates **5a** and **5b** was moderate (as in the control plate), with the exception of *Staphylococcus epidermidis*, whose growth was inhibited. In the cases where growth of the Gram-positive microorganisms was apparent, the production of fluorescent colonies was sometimes observed, but the intensity of fluorescence was weak compared to that produced by the Gram-negative microorganisms. Substrate **5c** was inhibitory to the growth of all of the Gram-positive microorganisms as well as the yeasts, hence no fluorescent colonies were generated. Thus, only the Gram-negative microorganisms produced fluorescence with substrate **5c** and this is clearly seen in the photograph of the agar plate shown in Figure 3.

		<b>5a</b>		<b>5b</b>		<b>5c</b>	
	<b>Microorganism / Reference<sup>a</sup></b>	<b>Growth<sup>b</sup></b>	<b>Fluorescence<sup>c</sup></b>	<b>Growth<sup>b</sup></b>	<b>Fluorescence<sup>c</sup></b>	<b>Growth<sup>b</sup></b>	<b>Fluorescence<sup>c</sup></b>
	Gram-negative microorganisms						
1	<i>Escherichia coli</i> NCTC 10418	++	++ blue	++	++ green	++	++ blue
2	<i>Klebsiella pneumoniae</i> NCTC 9528	++	++ blue	++	++ green	++	++ blue
3	<i>Providencia rettgeri</i> NCTC 7475	++	++ blue	++	++ green	++	++ blue
4	<i>Enterobacter cloacae</i> NCTC 11936	++	++ blue	++	++ green	++	++ blue
5	<i>Serratia marcescens</i> NCTC 10211	++	++ blue	++	++ green	++	++ blue
6	<i>Salmonella typhimurium</i> NCTC 74	++	++ blue	++	++ green	++	++ blue
7	<i>Pseudomonas aeruginosa</i> NCTC 10662	++	++ blue	++	++ green	++	+ blue
8	<i>Yersinia enterocolitica</i> NCTC 11176	++	++ blue	++	++ green	++	++ blue
9	<i>Burkholderia cepacia</i> NCTC 10743	++	+/- blue	++	+ green	++	++ blue
10	<i>Acinetobacter baumannii</i> NCTC 12156	++	++ blue	++	++ green	++	++ blue
	Gram-positive microorganisms						
11	<i>Streptococcus pyogenes</i> NCTC 8306	+	-	+	Tr green	-	-
12	<i>Staphylococcus aureus</i> (MRSA) NCTC 11939	+	-	+	-	-	-
13	<i>Staphylococcus aureus</i> (MSSA) NCTC 6571	+	-	+	-	-	-
14	<i>Staphylococcus epidermidis</i> NCTC 11047	-	-	-	-	-	-
15	<i>Listeria monocytogenes</i> NCTC 11994	+	Tr blue	+	Tr green	-	-
16	<i>Enterococcus faecium</i> NCTC 7171	+	Tr blue	+	+ green	-	-
17	<i>Enterococcus faecalis</i> NCTC 775	+	Tr blue	+	+ green	-	-
18	<i>Bacillus subtilis</i> ATCC 9372	+	-	+	-	-	-


	Yeasts						
19	<i>Candida albicans</i> ATCC 90028	+	Tr blue	+	-	-	-
20	<i>Candida glabrata</i> NCPF 3943	+	-	+	-	-	-

<sup>a</sup>NCTC: National Collection of Type Cultures; ATCC: American Type Culture Collection; NCPF: National Collection of Pathogenic Fungi.

<sup>b</sup>++ strong growth, + moderate growth, +/- weak growth.

<sup>c</sup>++ strong fluorescence, + moderate fluorescence, +/- weak fluorescence, Tr trace of fluorescence.

**Table 1.** Evaluation of the substrates **5a-c**. Substrate concentration = 100 mg L<sup>-1</sup>; inoculum = 100 000 colony-forming units (cfu)/spot.



**Figure 3.** An illustrative Columbia agar plate depicting the fluorescence generated by various microorganisms with substrate **5c**. <sup>a</sup> Microorganisms are numbered in the sequence shown in the Tables. Green spots represent Gram-negative bacteria, blue spots represent Gram-positive bacteria and purple spots represent the yeast species. Plate viewed under UV light ( $\lambda = 365$  nm).

Good growth of the Gram-negative microorganisms in the presence of substrates **6-8** was observed and moderate growth was noted for the majority of the Gram-positive microorganisms and the yeasts (Table 2). All of the Gram-negative microorganisms produced fluorescent colonies with substrates **6-8** with substrate **6** giving strong fluorescence and substrates **7** and **8** showing moderately intense fluorescence. Only substrate **6** generated moderately intense, fluorescent colonies with some of the Gram-positive microorganisms. There is an interesting contrast between the quinoline-derived substrate **5c** and the coumarin-derived substrates **7** and **8**. Only the Gram-negative microorganisms produced fluorescent colonies with these three substrates; however



substrate **5c** was inhibitory to the growth of Gram-positive microorganisms whereas substrates **7** and **8** were not inhibitory to these microorganisms. Thus, substrates **7** and **8** could potentially be used for differentiation of Gram-negative and Gram-positive microorganisms. Alternatively, for applications that target Gram-negative bacteria specifically, substrate **5c** could potentially be used as a both a fluorogenic indicator and a selective agent for inhibition of Gram-positive bacteria.

A yellow precipitate was noted in the agar plate containing substrate **6**; presumably this substrate has limited solubility in agar media at a concentration of 100 mg L<sup>-1</sup>. All of the three coumarin substrates **6-8** were therefore evaluated at a concentration of 25 mg L<sup>-1</sup> and this resulted in broadly similar growth and fluorescence patterns compared to the higher substrate concentration. However, even with this lower substrate concentration, a yellow precipitate was still apparent with substrate **6**. The concentration of the substrate **6** was therefore reduced further to 10 mg L<sup>-1</sup>. At this substrate concentration, the observed growth and fluorescence was again broadly similar to that at higher concentrations but no substrate precipitation was noted.

		<b>6</b>		<b>7</b>		<b>8</b>	
	<b>Microorganism / Reference<sup>a</sup></b>	<b>Growth<sup>b</sup></b>	<b>Fluorescence<sup>c</sup></b>	<b>Growth<sup>b</sup></b>	<b>Fluorescence<sup>c</sup></b>	<b>Growth<sup>b</sup></b>	<b>Fluorescence<sup>c</sup></b>
	Gram-negative microorganisms						
1	<i>Escherichia coli</i> NCTC 10418	++	++ yellow	++	+ green	++	+ yellow
2	<i>Klebsiella pneumoniae</i> NCTC 9528	++	++ yellow	++	+ green	++	+ yellow
3	<i>Providencia rettgeri</i> NCTC 7475	++	++ yellow	++	+ green	++	+ yellow
4	<i>Enterobacter cloacae</i> NCTC 11936	++	++ yellow	++	+ green	++	+ yellow
5	<i>Serratia marcescens</i> NCTC 10211	++	++ yellow	++	+ green	++	+ yellow
6	<i>Salmonella typhimurium</i> NCTC 74	++	++ yellow	++	+ green	++	+ yellow
7	<i>Pseudomonas aeruginosa</i> NCTC 10662	++	++ yellow	++	+ green	++	+ yellow
8	<i>Yersinia enterocolitica</i> NCTC 11176	++	+ yellow	++	+ green	++	+ yellow
9	<i>Burkholderia cepacia</i> NCTC 10743	++	++ yellow	++	+ green	++	+/- yellow
10	<i>Acinetobacter baumannii</i> NCTC 12156	++	++ yellow	++	+ green	++	+ yellow

	Gram-positive microorganisms						
11	<i>Streptococcus pyogenes</i> NCTC 8306	+	-	+	-	+	-
12	<i>Staphylococcus aureus</i> (MRSA) NCTC 11939	+	+ blue	+	-	+	-
13	<i>Staphylococcus aureus</i> (MSSA) NCTC 6571	+	+ blue	+	-	+	-
14	<i>Staphylococcus epidermidis</i> NCTC 11047	-	-	+	-	+	-
15	<i>Listeria monocytogenes</i> NCTC 11994	+	+ blue	+	-	+	-
16	<i>Enterococcus faecium</i> NCTC 7171	+	+ yellow	+	-	+	-
17	<i>Enterococcus faecalis</i> NCTC 775	+	+ yellow	+	-	+	-
18	<i>Bacillus subtilis</i> ATCC 9372	-	-	+	-	+	-
	Yeasts						
19	<i>Candida albicans</i> ATCC 90028	+	-	+	-	+	-
20	<i>Candida glabrata</i> NCPF 3943	-	-	+	-	+	-

<sup>a</sup>NCTC: National Collection of Type Cultures; ATCC: American Type Culture Collection; NCPF: National Collection of Pathogenic Fungi.

<sup>b</sup> ++ strong growth, + moderate growth, +/- weak growth.

<sup>c</sup> ++ strong fluorescence, + moderate fluorescence, +/- weak fluorescence.

**Table 2.** Evaluation of the substrates **6-8**. Substrate concentration = 100 mg L<sup>-1</sup>; inoculum = 100 000 colony-forming units (cfu)/spot.

In conclusion, all of the six novel fluorogenic substrates described in this paper produced fluorescent colonies with Gram-negative microorganisms in Columbia agar media. The growth of the Gram-negative microorganisms was not inhibited by these substrates. Moderate growth of the majority of the Gram-positive microorganisms and yeasts occurred in the presence of substrates **5a**, **5b** and **6** and fluorescent colonies were observed, but not with all of these microorganisms. Moderate growth of all of the Gram-positive microorganisms and yeast also occurred with substrates **7** and **8** but no fluorescent colonies were produced, thus allowing differentiation between these microorganisms

and the Gram-negative microorganisms. Substrate **5c** was inhibitory to the Gram-positive microorganisms and the yeasts thus creating a clear distinction between these microorganisms and the Gram-negative microorganisms.

## Acknowledgements

We thank bioMérieux SA for generous financial support and the National Mass Spectrometry Facility (NMSF) at Swansea University for high resolution mass spectra.

## References

1. Carlone GM, Valadez MJ, Pickett MJ. Methods for distinguishing Gram-positive from Gram-negative bacteria. *J Clin Microbiol.* 1982;16:1157-1159.
2. Anderson RJ, Groundwater PW, Huang Y, James AL, Orega S, Rigby A, Roger-Dalbert C, Perry JD. Synthesis and evaluation of novel chromogenic peptidase substrates based on 9-(4'-aminophenyl)-10-methylacridinium salts as diagnostic tools in clinical bacteriology. *Bioorg Med Chem Lett.* 2008;18:832-835.
3. Cellier M, Fabrega OL, Fazackerley E, James AL, Orega S, Perry JD, Salwatura VL, Stanforth SP. 2-Arylbenzothiazole, benzoxazole and benzimidazole derivatives as fluorogenic substrates for the detection of nitroreductase and aminopeptidase activity in clinically important bacteria. *Bioorg Med Chem.* 2011;19:2903-2910.
4. Cellier M, Fazackerley E, James AL, Orega S, Perry JD, Turnbull G, Stanforth SP. Synthesis of 2-arylbenzothiazole derivatives and their application in bacterial detection. Synthesis of 2-arylbenzothiazole derivatives and their application in bacterial detection. *Bioorg Med Chem.* 2014;22:1250-1261.
5. Dong X, Heo CH, Chen S, Kim HM, Liu Z. Quinoline-based two-photon fluorescent probe for nitric oxide in live cells and tissues. *Anal Chem.* 2014;86:308-311.
6. For example: Kim HJ, Han JH, Kim MK, Lim CS, Kim HM, Cho BR. Dual-color imaging of sodium/calcium ion activities with two-photon fluorescent probes. *Angew Chem Int Ed.* 2010;49:6786-6789; Han JH, Park SK, Lim CS, Park MK, Kim HJ, Kim HM, Cho BR. Simultaneous imaging of mitochondria and lysosomes by using two-photon fluorescent probes. *Chem Eur J.* 2012;18:15246-15249; Kim HJ, Heo CH, Kim HM. Benzimidazole-based Ratiometric two-photon fluorescent probes for acidic pH in live cells and tissues. *J Am Chem Soc.* 2013;135:17969-17977.
7. Manafi M, Kneifel W, Bascomb S. Fluorogenic and chromogenic substrates used in bacterial diagnostics. *Microbiol Rev.*, 1991;55:335-348 and references therein; Manafi M, Kneifel W. Rapid methods for differentiating Gram-positive from Gram-negative aerobic and facultative anaerobic bacteria. *J Appl Bacteriol.* 1990;69:822-827.
8. Agban A, Ounanian M, Luu-Duc C, Monget D. Synthesis of new fluorogenic substrate derivatives of 7-amino-4-trifluoromethylcoumarin. Detection of Gram-negative bacteria, Streptococci group A and Enterococci. *Ann. Pharm. Fr.* 1990;48:326-334.
9. Bhusal RP, Cho PY, Kim S-A, Park, H, Kim HS. Synthesis of green emitting coumarin bioconjugate for the selective determination of flu antigen. *Bull Korean Chem Soc.* 2011;32:1461-1462.
10. Tavares FX, Al-Barazanji KA, Bigham EC, Bishop MJ, Britt CS, Carlton DL, Feldman PL, Goetz AS, Grizzle MK, Guo YC, Handlon AL, Hertzog DL, Ignar DM, Lang DG, Ott RJ, Peat AJ, Zhou H-Q. Potent, selective, and orally efficacious antagonists of melanin-concentrating hormone receptor 1. *J Med Chem.* 2006;49:7095-7107.

11. Hofsløkken NU, Skattebøl L. Convenient method for the *ortho*-formylation of phenols. *Acta Chem Scand.* 1999;53:258-262.